

WAYS HEALTHCARE PROVIDERS CAN PREVENT OVERDOSE

6 SIMPLE STEPS FOR LIFESAVING IMPACT

STEP 1

Ask about overdose

Experiencing previous overdose is the greatest risk factor for subsequent overdose.^{1,2}

STEP 2

Prescribe buprenorphine

Individuals on MOUD have a lower risk of overdose and overdose-related mortality.³

STEP 3

Prescribe naloxone with every encounter

High coverage of naloxone is associated with a reduction in overdose.⁴

STEP 4

Watch for abstinence

Recent release from abstinence-based treatment or incarceration are associated with increased rates of overdose mortality.⁵

STEP 5





Screen for polysubstance use & avoid benzodiazepines

Polysubstance use, especially benzodiazepines, are associated with increased rates of overdose mortality.⁶

STEP 6

Get educated on overdose prevention sites

Overdose prevention sites have been associated with reduction in overdose mortality.⁷



READY TO TAKE ACTION?

Our team of addiction medicine experts are here to help.

855-337-MACS
MARYLANDMACS.ORG



MACS
Maryland Addiction Consultation Service

1. Britton PC, Wines JD Jr, Conner KR. Non-fatal overdose in the 12 months following treatment for substance use disorders. *Drug Alcohol Depend.* 2010 Feb 1;107(1):51-5. doi: 10.1016/j.drugalcdep.2009.09.005. Epub 2009 Oct 13. PMID: 19828263; PMCID: PMC2818271.
2. Coffin PO, Tracy M, Bucciarelli A, Ompad D, Vlahov D, Galea S. Identifying injection drug users at risk of nonfatal overdose. *Acad Emerg Med.* 2007 Jul;14(7):616-23. doi: 10.1197/j.aem.2007.04.005. Epub 2007 Jun 6. PMID: 17554010.
3. Wakeman SE, Larochelle MR, Ameli O, et al. Comparative Effectiveness of Different Treatment Pathways for Opioid Use Disorder. *JAMA Netw Open.* 2020;3(2):e1920622. doi:10.1001/jamanetworkopen.2019.20622
4. Walley AY, Xuan Z, Hackman HR, Quinn E, Doe-Simkins M, Sorensen-Alawad A, Ruiz S, Ozonoff A. Opioid overdose rates and implementation of naloxone distribution in Massachusetts: interrupted time series analysis. *BMJ.* 2013 Jan 30;346:f174. doi: 10.1136/bmj.f174. PMID: 23372174; PMCID: PMC4688551.
5. Waddell EN, Baker R, Hartung DM, et al. Reducing overdose after release from incarceration (ROAR): study protocol for an intervention to reduce risk of fatal and non-fatal opioid overdose among women after release from prison. *Health Justice.* 2020;8(1):18. Published 2020 Jul 10. doi:10.1186/s40352-020-00113-7
6. Compton WM, Valentino RJ, DuPont RL. Polysubstance use in the U.S. opioid crisis. *Molecular Psychiatry.* 2020 26:1. 2020;26(1):41-50. doi:10.1038/s41380-020-00949-3
7. Finke J, Chan J. The Case for Supervised Injection Sites in the United States. *American Family Physician.* 2022;105(5):454-455. Accessed August 17, 2022. <https://www.aafp.org/pubs/afp/issues/2022/0500/p454.html>